

STONE SPECTRA OF VON NEUMANN ALGEBRAS OF TYPE I_n

ANDREAS DÖRING
IAMPH, FACHBEREICH MATHEMATIK
J. W. GOETHE-UNIVERSITÄT FRANKFURT, GERMANY

ABSTRACT. The Stone spectrum of a von Neumann algebra is a generalization of the Gelfand spectrum, as was shown by de Groote. In this article we clarify the structure of the Stone spectra of von Neumann algebras of type I_n .

1. INTRODUCTION

In a new approach tying together classical and quantum observables, de Groote has developed the theory of *Stone spectra* of lattices and *observable functions*. In this section, we will give the relevant definitions and cite some of de Groote's results. For details, the reader is referred to de Groote's work [deG01, deG05].

When speaking of a lattice, we will always mean a σ -complete lattice at least. A lattice $\mathbb{L}$ always has a zero element 0 and a unit element 1. The starting point is the new notion of a *quasipoint* of a lattice, which is nothing but a maximal filter base, generalizing to arbitrary lattices what Stone did in the 1930s for Boolean algebras [Sto36]:

Definition 1. A subset $\mathfrak{B}$ of a lattice $\mathbb{L}$ is called a **quasipoint of $\mathbb{L}$** if it has the following properties:

- (i) $0 \notin \mathfrak{B}$,
- (ii) $\forall a, b \in \mathfrak{B} \exists c \in \mathfrak{B} : c \leq a \wedge b$,
- (iii) $\mathfrak{B}$ is maximal with respect to (i) and (ii).

It is easily seen that for a quasipoint $\mathfrak{B}$ of the lattice $\mathbb{L}$, we have

$$\forall a \in \mathfrak{B} \forall b \in \mathbb{L} : (a \leq b \implies b \in \mathfrak{B}).$$

In particular, $\forall a, b \in \mathfrak{B} : a \wedge b \in \mathfrak{B}$, so quasipoints are maximal dual ideals also [Bir73]. The set of quasipoints of a lattice $\mathbb{L}$ is denoted by $\mathcal{Q}(\mathbb{L})$ and is equipped with a natural topology (also inspired by Stone): for $a \in \mathbb{L}$, let

$$\mathcal{Q}_a(\mathbb{L}) := \{\mathfrak{B} \in \mathcal{Q}(\mathbb{L}) \mid a \in \mathfrak{B}\}.$$

Obviously, we have

$$\mathcal{Q}_{a \wedge b}(\mathbb{L}) = \mathcal{Q}_a(\mathbb{L}) \cap \mathcal{Q}_b(\mathbb{L}),$$

so $\{\mathcal{Q}_a(\mathbb{L}) \mid a \in \mathbb{L}\}$ is the base of a topology on $\mathbb{L}$.

Definition 2. The set $\mathcal{Q}(\mathbb{L})$ of quasipoints of a lattice $\mathbb{L}$, equipped with the topology given by the sets $\mathcal{Q}_a(\mathbb{L})$ defined above, is called the **Stone spectrum of the lattice $\mathbb{L}$** .

One can show that the Stone spectrum $\mathcal{Q}(\mathbb{L})$ is a zero-dimensional, completely regular Hausdorff space. For the example $\mathbb{L} = \mathbb{L}(\mathcal{H})$, the lattice of closed subspaces of a complex separable Hilbert space $\mathcal{H}$, $\mathcal{Q}(\mathbb{L})$ is not compact if $\dim \mathcal{H} > 1$. If $\mathcal{H}$ is infinite-dimensional, then $\mathcal{Q}(\mathbb{L})$ is not even locally compact [deG01]. The lattice $\mathbb{L}(\mathcal{H})$ plays an important role in the foundations of quantum theory, which was first recognized by Birkhoff and von Neumann [BirvNeu36]. $\mathbb{L}(\mathcal{H})$ is isomorphic to the $\mathcal{P}(\mathcal{H})$, the lattice of projections onto closed subspaces of $\mathcal{H}$.

Let $\mathcal{R}$ be a unital von Neumann algebra, given as a subalgebra of the algebra $\mathcal{L}(\mathcal{H})$ of bounded operators on some Hilbert space $\mathcal{H}$. The Stone spectrum $\mathcal{Q}(\mathcal{R})$ of $\mathcal{R}$ means the Stone spectrum $\mathcal{Q}(\mathcal{P}(\mathcal{R}))$ of the projection lattice $\mathcal{P}(\mathcal{R})$ of $\mathcal{R}$. In quantum theory, including quantum mechanics in the von Neumann representation, quantum field theory and quantum information theory, *observables* are represented by self-adjoint operators A in some von Neumann algebra $\mathcal{R}$, the *algebra of observables*. The set of observables $\mathcal{R}_{sa}$ forms a real linear space in the algebra $\mathcal{R}$.

De Groote shows that if $\mathcal{R}$ is abelian, there is a homeomorphism between the Stone spectrum $\mathcal{Q}(\mathcal{R}) = \mathcal{Q}(\mathcal{P}(\mathcal{R}))$ and the Gelfand spectrum $\Omega(\mathcal{R})$ of $\mathcal{R}$. Hence, for an arbitrary von Neumann algebra $\mathcal{R}$, the Stone spectrum $\mathcal{Q}(\mathcal{R})$ is a *generalization of the Gelfand spectrum*.

Observable functions are introduced in the following way:

Definition 3. Let $A \in \mathcal{R}_{sa}$, and let $E^A = (E_\lambda^A)_{\lambda \in \mathbb{R}}$ be the spectral family of A . The function

$$f_A : \mathcal{Q}(\mathcal{R}) \longrightarrow \mathbb{R}$$

defined by

$$f_A(\mathfrak{B}) := \inf\{\lambda \in \mathbb{R} \mid E_\lambda^A \in \mathfrak{B}\}$$

is called the **observable function corresponding to A** . $Ob(\mathcal{R}) := \{f_A \mid A \in \mathcal{R}_{sa}\}$ denotes the set of observable functions of $\mathcal{R}$.

One can show that the image of f_A is the spectrum of A . Observable functions are continuous functions, so

$$Ob(\mathcal{R}) \subseteq C_b(\mathcal{Q}(\mathcal{R}), \mathbb{R}),$$

where $C_b(\mathcal{Q}(\mathcal{R}), \mathbb{R})$ denotes the set of continuous bounded real-valued functions on the Stone spectrum $\mathcal{Q}(\mathcal{R})$ of $\mathcal{R}$. De Groote shows that equality only holds for abelian von Neumann algebras $\mathcal{R}$. Moreover, if $\mathcal{R}$ is abelian, then the observable functions turn out to be the Gelfand transforms of the self-adjoint elements of $\mathcal{R}$, so for an arbitrary von Neumann algebra, the observable functions are *generalized Gelfand transforms*.

There are many more results on Stone spectra and observable functions and their relation to physics [deG01, deG05]. For example, observable functions can be characterized intrinsically, without reference to self-adjoint operators. Stone spectra also play some role in the proof of the generalized Kochen-Specker theorem, which is an important no-go theorem on hidden variables in quantum theory [KocSpe67, Doe04].

The elements of the Stone spectrum $\mathcal{Q}(\mathcal{R})$, the quasipoints, are defined using Zorn's lemma. As usual, it is not easy to get some intuition of such objects. Some extra structure of the von Neumann algebra is needed to clarify the properties of the quasipoints and the Stone spectrum. The quasipoints and the Stone spectrum of type- I_n -factors $\mathcal{R} = \mathcal{L}(\mathcal{H})$, where $\dim \mathcal{H} = n \in \{0, 1, \dots\}$ is finite, are known [deG01]. We need the following: an isolated point $\mathfrak{B}$ of the Stone spectrum $\mathcal{Q}(\mathcal{R})$ is called an *atomic quasipoint*. If $\mathcal{R} = \mathcal{L}(\mathcal{H})$, then the atomic quasipoints of $\mathcal{P}(\mathcal{R}) (= \mathcal{P}(\mathcal{H}))$ are of the form

$$\mathfrak{B}_{\mathbb{C}x} = \{P \in \mathcal{P}(\mathcal{H}) \mid P \geq P_{\mathbb{C}x}\},$$

where $x \in \mathcal{H}$, $|x| = 1$ and $P_{\mathbb{C}x}$ is the projection onto the line $\mathbb{C}x$. While for infinite-dimensional $\mathcal{H}$, $\mathcal{P}(\mathcal{H})$ also has non-atomic quasipoints, for finite-dimensional $\mathcal{H}$ the situation is simple:

Proposition 4. *If $\mathcal{H}$ is finite-dimensional, there are only atomic quasipoints in $\mathcal{P}(\mathcal{H})$, and we have*

$$\mathcal{Q}(\mathcal{P}(\mathcal{H})) = \{\mathfrak{B}_{\mathbb{C}x} \mid x \in S^1(\mathcal{H})\},$$

where $S^1(\mathcal{H})$ denotes the unit sphere in Hilbert space. For finite-dimensional $\mathcal{H}$, $\mathcal{P}(\mathcal{H}) \simeq \mathbb{L}(\mathcal{H})$ is the projection lattice $\mathcal{P}(\mathcal{R})$ of a type I_n factor $\mathcal{R}$, where $n = \dim \mathcal{H}$. This type I_n factor simply is represented as $\mathbb{M}_n(\mathbb{C})$, the $n \times n$ complex matrices acting on $\mathcal{H}$.

Of course, the inner product of $\mathcal{H}$ plays no role here, but only the linear structure, so we have also characterized the quasipoints of the lattice of subspaces of a finite-dimensional vector space.

There were no results on the structure of quasipoints and Stone spectra of more general von Neumann algebras up to now. In this article, we will examine von Neumann algebras of type I_n for finite n . Type I_n algebras include all von Neumann algebras on finite-dimensional Hilbert spaces and all abelian von Neumann algebras. (The latter are of those of type I_1 .) We will make use of the fact that such an algebra is of the form $\mathbb{M}_n(\mathcal{A})$, where $\mathcal{A}$ is the center of $\mathcal{R}$. Note that while $\mathcal{R} \simeq \mathbb{M}_n(\mathcal{A})$ is given by “finite” $n \times n$ -matrices, the center $\mathcal{A}$ of $\mathcal{R}$ may be represented on an infinite-dimensional Hilbert space.

Drawing on a result on *abelian quasipoints* (section 2), i.e. quasipoints containing an abelian projection, a fairly complete characterization of the Stone spectrum of a type I_n algebra is obtained. In particular, all quasipoints of a type I_n algebra are abelian (Thm. 34) and the orbits of the action of the unitary group on the Stone spectrum $\mathcal{Q}(\mathcal{R})$ are parametrized by the quasipoints of the center of $\mathcal{R}$ (Thm. 36).

2. ABELIAN QUASIPOINTS OF VON NEUMANN ALGEBRAS

In this section, we will regard quasipoints containing an abelian projection. It will be shown that there is a close relationship between the abelian quasipoints of a von Neumann algebra $\mathcal{R}$ and the quasipoints of the center of $\mathcal{R}$. This result will be central to the classification of Stone spectra of type I_n von Neumann algebras.

Definition 5. *A quasipoint $\mathfrak{B} \subseteq \mathcal{P}(\mathcal{R})$ is called **abelian** if it contains an abelian projection $E \in \mathcal{R}$. The set of abelian quasipoints of a von Neumann algebra $\mathcal{R}$ is denoted by $\mathcal{Q}^{ab}(\mathcal{R})$.*

Definition 6. The *E-trunk* $\mathfrak{B}_E$ ($E \in \mathfrak{B}$) of a quasipoint $\mathfrak{B}$ is the set

$$\mathfrak{B}_E := \{F \in \mathfrak{B} \mid F \leq E\}.$$

Obviously, $\mathfrak{B}_E$ is a filter basis.

Lemma 7. The *E-trunk* $\mathfrak{B}_E$ uniquely determines the quasipoint $\mathfrak{B}$.

Proof. Let $\mathfrak{B}_1, \mathfrak{B}_2$ be two quasipoints whose *E-trunk* is $\mathfrak{B}_E$. Let F be a projection in $\mathfrak{B}_1$. Then we have $E \wedge F \in \mathfrak{B}_E \subset \mathfrak{B}_2$. If a quasipoint contains a projection, it contains all larger projections, so $F \in \mathfrak{B}_2$ and $\mathfrak{B}_1 = \mathfrak{B}_2$ follows. $\square$

This lemma holds analogously for any lattice $\mathbb{L}$, since no features of the von Neumann algebra are used.

Definition 8. Let $\mathcal{R} \subseteq \mathcal{L}(\mathcal{H})$ be a von Neumann algebra, $\mathfrak{B} \subset \mathcal{Q}_E(\mathcal{R})$ a quasipoint containing E and $\theta \in \mathcal{R}$ a partial isometry such that $E = \theta^*\theta$. We set

$$\theta(\mathfrak{B}_E) := \{\theta F \theta^* \mid F \in \mathfrak{B}_E\}.$$

Lemma 9. If $\mathcal{R} \subseteq \mathcal{L}(\mathcal{H})$ is a von Neumann algebra and $\theta \in \mathcal{R}$ is a partial isometry such that $E := \theta^*\theta$, then for all projections $P_U \in \mathcal{R}$ such that $P_U \leq E$ it holds that

$$\theta P_U \theta^* = P_{\theta U}.$$

Proof. For $x \in U$, we have

$$\theta P_U \theta^* \theta x = \theta P_U E x = \theta x = P_{\theta U} \theta x.$$

If $y \in (\theta U)^\perp$, then $\theta^* y \in U^\perp$ and thus

$$\theta P_U \theta^* y = 0 = P_{\theta U} y.$$

$\square$

Obviously, $\theta(\mathfrak{B}_E)$ is the $\theta E \theta^*$ -trunk of a quasipoint of $\mathcal{R}$ (given that $\theta^*\theta = E$). We will denote the quasipoint induced by $\theta E \theta^*$ by $\theta_Q(\mathfrak{B}_E)$.

Remark 10. Since in general $\theta^*\theta \notin \mathfrak{B}$ for an arbitrary partial isometry θ and an arbitrary quasipoint $\mathfrak{B}$, we have no action of the set of partial isometries on the Stone spectrum $\mathcal{Q}(\mathcal{R})$. On the other hand, if θ is unitary, we can define an operation, see subsection 3.

Let $\mathfrak{B} \in \mathcal{Q}^{ab}(\mathcal{R})$ be an abelian quasipoint, and let $E \in \mathfrak{B}$ be an abelian projection. Each $F \in \mathfrak{B}_E$ is a subprojection of the abelian projection E and hence of the form $F = QE$, where $Q \in \mathcal{R}$ is a central projection. Then $Q \in \mathfrak{B}$ holds, so $Q \in \mathfrak{B} \cap \mathcal{C}$. If, conversely, $Q \in \mathfrak{B} \cap \mathcal{C}$ holds, then $QE \in \mathfrak{B}_E$, therefore we have

$$\mathfrak{B}_E = \{QE \mid Q \in \mathfrak{B} \cap \mathcal{C}\}.$$

The mapping

$$\begin{aligned} \zeta_E : \mathcal{R}'E &\longrightarrow \mathcal{R}'C_E, \\ T'E &\longmapsto T'C_E \end{aligned}$$

is a $*$ -isomorphism (see Prop. 5.5.5 in [KadRinI97]). Since

$$\begin{aligned} PE \wedge QE &= PQE = (P \wedge Q)E, \\ PE \vee QE &= (P + Q)E - PQE = (P \vee Q)E, \end{aligned}$$

$\zeta_E|_{\mathfrak{B}_E}$ is a lattice isomorphism from $\mathfrak{B}_E$ onto $(\mathfrak{B} \cap \mathcal{C})C_E$. $\mathfrak{B} \cap \mathcal{C}$ is a quasipoint of $\mathcal{P}(\mathcal{C})$: obviously, $\mathfrak{B} \cap \mathcal{C}$ is a filter basis $\mathcal{P}(\mathcal{C})$. Let β be a quasipoint in $\mathcal{P}(\mathcal{C})$ containing $\mathfrak{B} \cap \mathcal{C}$. Assume that $P \in \beta \setminus (\mathfrak{B} \cap \mathcal{C})$. Then $PC_E \in \beta \setminus (\mathfrak{B} \cap \mathcal{C})$ holds, therefore $PE \notin \mathfrak{B}_E$. Then there is some $QE \in \mathfrak{B}_E$ such that $PQE = 0$, so $PC_EQC_E = PQC_E = 0$, but that contradicts $PC_E, QC_E \in \beta$. Thus $\mathfrak{B} \cap \mathcal{C}$ is a quasipoint in $\mathcal{P}(\mathcal{C})$, and hence

$$\zeta_E(\mathfrak{B}_E) = (\mathfrak{B} \cap \mathcal{C})C_E.$$

In this manner, each abelian quasipoint $\mathfrak{B} \in \mathcal{Q}^{ab}(\mathcal{R})$ is assigned a quasipoint $\beta(\mathfrak{B}) := \mathfrak{B} \cap \mathcal{C} \in \mathcal{Q}(\mathcal{C})$ of the center $\mathcal{C}$ of $\mathcal{R}$. Moreover, the mapping

$$\begin{aligned} \zeta : \mathcal{Q}(\mathcal{R}) &\longrightarrow \mathcal{Q}(\mathcal{C}) \\ \mathfrak{B} &\longmapsto \mathfrak{B} \cap \mathcal{C} \end{aligned}$$

is surjective, since each quasipoint $\beta \in \mathcal{Q}(\mathcal{C})$ (being a filter base in $\mathcal{P}(\mathcal{R})$) is contained in some quasipoint $\mathfrak{B} \in \mathcal{Q}(\mathcal{R})$.

Let $\mathfrak{B}, \tilde{\mathfrak{B}} \in \mathcal{Q}^{ab}(\mathcal{R})$ be abelian quasipoints such that

$$\beta := \mathfrak{B} \cap \mathcal{C} = \tilde{\mathfrak{B}} \cap \mathcal{C}.$$

Let $E \in \mathfrak{B}, \tilde{E} \in \tilde{\mathfrak{B}}$ be abelian projections. Since $C_E, C_{\tilde{E}} \in \beta$, $C_EC_{\tilde{E}} \in \beta$ holds and $C_EC_{\tilde{E}}E \in \mathfrak{B}, C_EC_{\tilde{E}}\tilde{E} \in \tilde{\mathfrak{B}}$ are abelian projections with the same central carrier $C_EC_{\tilde{E}}$. Hence, without loss of generality, one can assume $C_E = C_{\tilde{E}}$. It follows that E and $\tilde{E}$ are equivalent (see Prop. 6.4.6 in [KadRinII97]). Let $\theta \in \mathcal{R}$ be a partial isometry such that $\theta^*\theta = E, \theta\theta^* = \tilde{E}$, therefore $\theta E\theta^* = \tilde{E}$. It follows that

$$\begin{aligned} \theta\mathfrak{B}_E\theta^* &= \{\theta QE\theta^* \mid Q \in \beta\} = \{Q\theta E\theta^* \mid Q \in \beta\} \\ &= \{Q\tilde{E} \mid Q \in \beta\} = \mathfrak{B}_{\tilde{E}}, \end{aligned}$$

so

$$\theta_{\mathcal{Q}}(\mathfrak{B}) = \tilde{\mathfrak{B}}.$$

Conversely, let $\mathfrak{B}, \tilde{\mathfrak{B}}$ be abelian quasipoints, and let $\theta \in \mathcal{R}$ be a partial isometry such that $E := \theta^*\theta \in \mathfrak{B}, \tilde{E} := \theta\theta^* \in \tilde{\mathfrak{B}}$. From this, $\theta_{\mathcal{Q}}(\mathfrak{B}) = \tilde{\mathfrak{B}}$ as shown. Let $F \in \mathfrak{B}$ be abelian, $F \leq E$. Then θF is a partial isometry from $(\theta F)^*\theta F = FEF = F$ to $\theta F\theta^* \in \tilde{\mathfrak{B}}$. Since $\theta F\theta^*$ is abelian, too, we can assume without loss of generality that E and $\tilde{E}$ are abelian. From the definition of $\theta_{\mathcal{Q}}$, it follows that

$$\tilde{\mathfrak{B}}_{\tilde{E}} = \theta\mathfrak{B}_E\theta^* = \{\theta QE\theta^* \mid Q \in \mathfrak{B} \cap \mathcal{C}\} = \{Q\tilde{E} \mid Q \in \mathfrak{B} \cap \mathcal{C}\}$$

holds, so

$$\{P\tilde{E} \mid P \in \tilde{\mathfrak{B}} \cap \mathcal{C}\} = \{Q\tilde{E} \mid Q \in \mathfrak{B} \cap \mathcal{C}\},$$

and hence, since $C_E = C_{\tilde{E}}$,

$$\begin{aligned} \{PC_E \mid P \in \tilde{\mathfrak{B}} \cap \mathcal{C}\} &= \{QC_E \mid Q \in \mathfrak{B} \cap \mathcal{C}\} \\ \iff (\tilde{\mathfrak{B}} \cap \mathcal{C})C_E &= (\mathfrak{B} \cap \mathcal{C})C_E, \end{aligned}$$

that is, $\tilde{\mathfrak{B}} \cap \mathcal{C} = \mathfrak{B} \cap \mathcal{C}$. Summing up, it is proven that:

Theorem 11. *Let $\mathcal{R}$ be a von Neumann algebra with center $\mathcal{C}$. Then the mapping*

$$\begin{aligned}\zeta : \mathcal{Q}(\mathcal{R}) &\longrightarrow \mathcal{Q}(\mathcal{C}) \\ \mathfrak{B} &\longmapsto \mathfrak{B} \cap \mathcal{C}\end{aligned}$$

is surjective. If $\mathfrak{B}, \tilde{\mathfrak{B}} \in \mathcal{Q}^{ab}(\mathcal{R})$ are two abelian quasipoints, then $\zeta(\mathfrak{B}) = \zeta(\tilde{\mathfrak{B}})$ holds if and only if there is a partial isometry $\theta \in \mathcal{R}$ with $\theta_{\mathcal{Q}}(\mathfrak{B}) = \tilde{\mathfrak{B}}$.

3. THE ACTION OF THE UNITARY GROUP ON THE STONE SPECTRUM $\mathcal{Q}(\mathcal{R})$

A unitary operator transforms a quasipoint in the obvious way:

Definition 12. *Let $T \in \mathcal{U}(\mathcal{H})$ be a unitary operator. T acts on $\mathfrak{B} \in \mathcal{Q}(\mathcal{R})$ ($\mathfrak{R} \subseteq \mathcal{L}(\mathcal{H})$) by*

$$T.\mathfrak{B} := \{TET^* \mid E \in \mathfrak{B}\}.$$

Lemma 13. *$T.\mathfrak{B}$ is a quasipoint of the von Neumann algebra $T\mathcal{R}T^*$. If $T \in \mathcal{U}(\mathcal{R})$, then $T.\mathfrak{B} \in \mathcal{Q}(\mathcal{R})$.*

Proof. We have

$$T(E \wedge F)T^* \leq TET^* \wedge TFT^*.$$

Moreover,

$$T^*(TET^* \wedge TFT^*)T \leq E \wedge F,$$

so $T(E \wedge F)T^* = TET^* \wedge TFT^*$. Thus $T.\mathfrak{B}$ is a filter basis and hence contained in some quasipoint $\mathfrak{B}' \in T\mathcal{R}T^*$. $T^*.\mathfrak{B}'$ also is a filterbasis. We have

$$T^*.(T.\mathfrak{B}) = \mathfrak{B} \subseteq T^*.\mathfrak{B}'.$$

From the maximality of $\mathfrak{B}$, equality holds. □

4. THE STONE SPECTRUM OF A TYPE I_n VON NEUMANN ALGEBRA

Let $\mathcal{R}$ be a von Neumann algebra of type I_n , n finite. We will show that every quasipoint $\mathfrak{B} \in \mathcal{Q}(\mathcal{R})$ of $\mathcal{R}$ is abelian, i.e. contains an abelian projection. In order to do so, we will use the fact that $\mathcal{R}$ is (isomorphic to) a $n \times n$ -matrix algebra, albeit with entries from another von Neumann algebra, the center of $\mathcal{R}$. We regard $\mathcal{R}$ as acting on the Hilbert module $\mathcal{A}^n$, which generalizes the vector space $\mathbb{C}^n$. The abelian projections will be those projecting onto “lines” of the form $a\mathcal{A}$, where $\mathcal{A} := \mathcal{C}(\mathcal{R})$ is the center of $\mathcal{R}$. Of course, $\mathcal{A}$ is not a field and $\mathcal{A}^n$ is not a vector space, so we cannot use arguments for subspace lattices of finite-dimensional vector spaces directly (in which case every quasipoint is abelian). But we will introduce equivalence relations on $\mathcal{A}$ and $\mathcal{A}^n$ that turn them into a field and an n -dimensional vector space, respectively, and show that after taking equivalence classes, enough of the structure remains intact to allow the conclusion that every quasipoint of $\mathcal{R}$ is abelian. The intuition from linear algebra carries through. From Thm. 11, we know that the abelian quasipoints can be mapped to quasipoints of the center of $\mathcal{R}$ via

$$\begin{aligned}\xi : \mathcal{Q}^{ab}(\mathcal{R}) &\longrightarrow \mathcal{Q}(\mathcal{C}), \\ \mathfrak{B} &\longmapsto \mathfrak{B} \cap \mathcal{C},\end{aligned}$$

where two quasipoints $\mathfrak{B}, \tilde{\mathfrak{B}}$ are mapped to the same quasipoint of the center if and only if there is a partial isometry $\theta \in \mathcal{R}$ such that $\theta_{\mathcal{Q}}(\mathfrak{B}) = \tilde{\mathfrak{B}}$. Using the fact

that $\mathcal{R}$ is a finite algebra, we can replace partial isometries with unitary operators. This will allow us to specify the orbits of the unitary group $\mathcal{U}(\mathcal{R})$ acting on $\mathcal{Q}(\mathcal{R})$ (Thm. 36).

4.1. Hilbert modules and the projections E_a . It is well known that each type I_n von Neumann algebra $\mathcal{R}$ is $*$ -isomorphic to $\mathbb{M}_n(\mathcal{A})$, the matrix algebra with entries from $\mathcal{A} = \mathcal{C}(\mathcal{R})$, the center of $\mathcal{R}$ (see Thm. 6.6.5 in [KadRinII97]). Let $\mathcal{A}^n$ be the free right module over $\mathcal{A}$ consisting of n copies of $\mathcal{A}$. Another common notation for $\mathbb{M}_n(\mathcal{A})$ is $\text{End}_{\mathcal{A}}(\mathcal{A}^n)$, the algebra of $\mathcal{A}$ -linear endomorphisms of $\mathcal{A}^n$. $\mathbb{M}_n(\mathcal{A})$ acts on the Hilbert space $\tilde{\mathcal{H}} := \bigoplus^n \mathcal{H}_{\mathcal{A}}$, the n -fold direct sum of $\mathcal{H}_{\mathcal{A}}$, which is the Hilbert space $\mathcal{A}$ acts on. We will not make use of $\tilde{\mathcal{H}}$ and the representation of $\mathbb{M}_n(\mathcal{A})$ on it, because we will regard $\mathbb{M}_n(\mathcal{A})$ as an algebra that acts on the $\mathcal{A}$ -module $\mathcal{A}^n$ from the left. Elements $a = (a_1 \oplus \dots \oplus a_n)^t$ of $\mathcal{A}^n$ are regarded as column vectors. The operation of $\mathbb{M}_n(\mathcal{A})$ on $\mathcal{A}^n$ is a “matrix $\times$ vector” operation. (Since $\mathcal{A}$ is commutative, $\mathcal{A}^n$ can be regarded as a left module as well. The chosen convention fits the natural structure of $\mathcal{A}^n$ as an $\mathbb{M}_n(\mathcal{A})$ - $\mathcal{A}$ -bimodule.)

$\mathcal{A}^n$ has a canonical basis with basis elements

$$e_j := (0 \oplus \dots \oplus 0 \oplus \overset{j}{\underset{\downarrow}{1}} \oplus 0 \oplus \dots \oplus 0)^t,$$

where 1 is the unit of $\mathcal{A}$. With respect to this basis, $a \in \mathcal{A}^n$ is denoted as $a = (a_1 \oplus \dots \oplus a_n)^t$. The sign of transposition will be omitted from now on.

There is an $\mathcal{A}$ -valued product defined on $\mathcal{A}^n$ such that $\mathcal{A}^n$ becomes a **Hilbert- $\mathcal{A}$ -module**. Since $\mathcal{A}^n$ is a right module, the inner product is $\mathcal{A}$ -linear with respect to the *second* variable:

$$\begin{aligned} (a|b) &= (a_1 \oplus \dots \oplus a_n | b_1 \oplus \dots \oplus b_n) := \sum_{k=1}^n a_k^* b_k, \\ (a|b\alpha) &= (a|b)\alpha = \alpha(a|b) = (a\alpha^*|b) \end{aligned}$$

for $a, b \in \mathcal{A}^n, \alpha \in \mathcal{A}$. In the second line the commutativity of $\mathcal{A}$ was employed. The inner product induces a norm on $\mathcal{A}^n$ by

$$|a| := |(a|a)|^{\frac{1}{2}},$$

where the norm on the right hand side is the norm on $\mathcal{A}$.

Let $\Omega := \mathcal{Q}(\mathcal{A})$ be the Stone spectrum of $\mathcal{A}$. Without loss of generality, we can assume $\mathcal{A} = C(\Omega)$. Let $(\Omega_1, \dots, \Omega_n)$ be a partition of Ω into closed-open sets $\Omega_k \neq \emptyset$, and let $a_k := \chi_{\Omega_k}$. Then for all $\beta \in \Omega$,

$$\sum_k |a_k(\beta)|^2 = 1$$

and hence $|a_1 \oplus \dots \oplus a_n| = 1$. Moreover, for $k \neq j$,

$$(a_k|a_j) = a_k^* a_j = 0,$$

that is, the a_k are pairwise orthogonal. But (the analogue of) Pythagoras' theorem does not hold, since for our example one obtains

$$|a_1 \oplus \dots \oplus a_n|^2 = 1 < n = \sum_{k=1}^n |a_k|^2.$$

In general, operators on Hilbert modules are –different from those on Hilbert spaces– not (all) adjointable, which is due to the lack of self-duality of Hilbert modules, see for example [WeO93, p 240]. A mapping $T : E \rightarrow E$ from a Hilbert module E to itself is called **adjointable** if there is a mapping T^* such that

$$(Ta|b) = (a|T^*b)$$

for all $a, b \in E$. The mapping T^* is called the **adjoint of T** . One can show that if T is adjointable, then T^* is unique, $T^{**} = T$ and both T and T^* are module maps which are bounded with respect to the operator norm ([WeO93, Lemma 15.2.3]).

For our purpose, we have to characterize the projections in $\mathcal{R} \simeq \mathbb{M}_n(\mathcal{A})$.

Lemma 14. *The elements of $\mathbb{M}_n(\mathcal{A})$ are adjointable, $T = T_{jk} \in \mathbb{M}_n(\mathcal{A})$ has adjoint $(T^*)_{jk} = T_{kj}^*$.*

Proof. Let $a, b \in \mathcal{A}^n, T = T_{jk} \in \mathbb{M}_n(\mathcal{A})$. We have

$$\begin{aligned} (Ta|b) &= \sum_{k,j} (T_{kj}a_j)^* b_k = \sum_{k,j} a_j^* T_{kj}^* b_k \\ &= \sum_{j,k} a_j^* (T_{jk})^* b_k = (a|T^*b). \end{aligned}$$

□

The adjoint of $T \in \mathbb{M}_n(\mathcal{A})$ in the Hilbert module sense thus is the usual, Hilbert space adjoint of T . It follows that the projections of the von Neumann algebra $\mathbb{M}_n(\mathcal{A})$ are the projections of the algebra $\mathbb{B}(\mathcal{A}^n)$ of adjointable operators of the Hilbert module $\mathcal{A}^n$. (A projection P is $\mathcal{A}$ -linear, so it is contained in $\mathbb{M}_n(\mathcal{A})$.)

We will now introduce projections E_a that map from $\mathcal{A}^n$ onto “lines” of the form $a\mathcal{A}$: let $a \in \mathcal{A}^n$ be such that $p := (a|a) \in \mathcal{A}$ is a projection. Then

$$\forall k \leq n : pa_k = a_k,$$

since for $\beta \in \mathcal{Q}(\mathcal{A})$ such that $a_k(\beta) \neq 0$, one obtains $p(\beta) = \sum_j a_j^* a_j \neq 0$ and hence $p(\beta) = 1$, since p is a projection. Here and in the following, the components $a_k \in \mathcal{A}$ of a are identified with their Gelfand transforms. We get

$$\forall \beta \in \mathcal{Q}(\mathcal{A}) \quad \forall k \leq n : (a_k p)(\beta) = a_k(\beta) p(\beta) = a_k(\beta),$$

so $ap = a$.

Now define

$$\begin{aligned} E_a : \mathcal{A}^n &\longrightarrow \mathcal{A}^n, \\ b &\longmapsto a(a|b). \end{aligned}$$

For all $b, c \in \mathcal{A}^n$,

$$\begin{aligned}(E_a b | c) &= (a(a|b) | c) = (a|b)^*(a|c) \\ &= (b|a)(a|c) = (b|a(a|c)) \\ &= (b|E_a c),\end{aligned}$$

so we have $E_a^* = E_a$. Moreover, for all $b \in \mathcal{A}^n$,

$$\begin{aligned}E_a^2 b &= a(a|E_a b) = a(a|a(a|b)) \\ &= a(a|a)(a|b) = ap(a|b) \\ &= a(a|b) = E_a b,\end{aligned}$$

so $E_a^2 = E_a$, that is, E_a is a projection with $\text{im } E_a \subseteq a\mathcal{A}$. In fact, equality holds: let $b = a\alpha \in a\mathcal{A}$. Then

$$a(a|b) = a(a|a)\alpha = a\alpha,$$

therefore $a\mathcal{A} \subseteq \text{im } E_a$. The central carrier C_{E_a} of E_a is $I_n(a|a)$: obviously, $I_n(a|a)$ is a central projection, since $I_n\mathcal{A} \simeq \mathcal{A}$ is the center of $\mathbb{M}_n(\mathcal{A})$. It holds that

$$\begin{aligned}E_a b(a|a) &= a(a|b)(a|a) = a(a|a)(a|b) \\ &= a(a|b) = E_a b,\end{aligned}$$

so $C_{E_a} \leq I_n(a|a)$. Conversely, let $I_n q$ be a central projection such that $qE_a = E_a q = E_a$. Then for all $b \in \mathcal{A}^n$,

$$\begin{aligned}E_a b &= a(a|b) = qE_a b = qa(a|b) \\ &= a(a|qb) = E_a(qb).\end{aligned}$$

In particular, one obtains

$$\begin{aligned}E_a(qa) &= E_a a \\ \iff a(a|qa) &= a(a|a) = ap = a \\ \iff a(a|a)q &= a \\ \iff aq = qa &= a \\ \implies q(a|a) &= qp = (a|a) = p \\ \implies p &\leq q,\end{aligned}$$

therefore, the central carrier of E_a is $C_{E_a} = I_n p = I_n(a|a)$.

The E_a are of interest, because they are *abelian* projections:

Lemma 15. *E_a is an abelian projection from $\mathcal{A}^n$ onto $a\mathcal{A}$ with central carrier $I_n(a|a)$.*

Proof. It only remains to show that E_a is abelian. Let $A, B \in \mathbb{M}_n(\mathcal{A})$. Then it holds for all $b \in \mathcal{A}^n$ that

$$\begin{aligned}E_a A E_a B E_a b &= E_a A E_a (Ba)(a|b) = E_a (Aa)(a|Ba)(a|b) \\ &= a(a|Aa)(a|Ba)(a|b) = a(a|Ba)(a|Aa)(a|b) \\ &= E_a (Ba)(a|Aa)(a|b) = E_a B E_a (Aa)(a|b) \\ &= E_a B E_a A E_a b,\end{aligned}$$

so $E_a A E_a B E_a = E_a B E_a A E_a$. □

E_a is a projection in $\mathbb{M}_n(\mathcal{A})$ if $(a|a)$ is a projection in $\mathcal{A}$. The converse is also true:

Remark 16. *Let $a \in \mathcal{A}^n$ be such that E_a is a projection. Then $(a|a) \in \mathcal{A}$ is a projection.*

Proof. From $E_a^2 = E_a$, $a(a|b) = a(a|E_a b) = a(a|a)(a|b)$ for all $b \in \mathcal{A}^n$. For $b = a$,

$$a(a|a) = a(a|a)^2.$$

This means that $(a|a) \in \{0, 1\}$ holds on the support

$$\text{supp } a := \bigcup_{k \leq n} \text{supp } a_k = \text{supp}(a|a).$$

If $\beta \in \Omega = \mathcal{Q}(\mathcal{A})$ is such that $(a|a)(\beta) \neq 0$, then $a_k(\beta) \neq 0$ holds for at least one $k \leq n$ and thus $(a|a)(\beta) = 1$. So $(a|a) = 1$ holds on $\text{supp}(a|a)$ and $(a|a)$ is a projection. $\square$

If $a_1, \dots, a_n \in \mathcal{A}$ are projections and $a := \sum_k a_k e_k$, then E_a is a projection if and only if the a_k are pairwise orthogonal, because, according to its definition and the above remark, E_a is a projection if and only if $(a|a)$ is a projection. For $a := \sum_k a_k e_k$, we have $(a|a) = \sum_k a_k$, and this is a projection if and only if the a_k are pairwise orthogonal. Furthermore, $\sum_k a_k E_{e_k} = \sum_k E_{a_k e_k}$ (the a_k are projections again): for all $b \in \mathcal{A}^n$, it holds that

$$\begin{aligned} \left(\sum_k a_k E_{e_k}\right)b &= \sum_k a_k e_k (e_k|b) = \sum_k a_k^2 e_k (e_k|b) \\ &= \sum_k a_k e_k (e_k|b a_k) = \sum_k a_k e_k (a_k e_k|b) \\ &= \sum_k E_{a_k e_k} b. \end{aligned}$$

Lemma 17. *$A := \sum_{k=1}^n a_k E_{e_k}$ is a projection if and only if all the a_k are projections. In this case, the central carrier of A is $C_A = I_n(\bigvee_k a_k)$.*

Proof. From $(e_j|e_k) = \delta_{jk} e_k$ we get

$$\forall c \in \mathcal{A}^n : E_{e_j} E_{e_k} c = e_j (e_j|E_{e_k} c) = e_j (e_j|e_k)(e_k|c) = \delta_{jk} E_j c,$$

so $E_{e_j} E_{e_k} = \delta_{jk} E_{e_j}$. $A := \sum_{k=1}^n a_k E_{e_k}$ is a projection if and only if $a_k^* = a_k$ holds for all $k \leq n$ and if $A^2 = A$. Since we have

$$\begin{aligned} A^2 &= \left(\sum_{k=1}^n a_k E_{e_k}\right)^2 = \sum_{j,k=1}^n a_j a_k E_{e_j} E_{e_k} \\ &= \sum_{j,k=1}^n a_j a_k \delta_{jk} E_{e_j} = \sum_{j=1}^n a_j^2 E_{e_j}, \end{aligned}$$

this holds if and only if all the $a_j \in \mathcal{A}$ are projections. $C_A = I_n(\bigvee_k a_k)$ holds then, obviously. $\square$

Remark 18. *With respect to the basis $(e_1, \dots, e_n)$ of $\mathcal{A}^n$, E_a has the matrix*

$$(E_a)_{jk} = (a_j a_k^*)_{j,k \leq n}.$$

Proof. It holds that

$$\begin{aligned} E_a e_k &= a(a|e_k) = a \sum_j a_j^* \delta_{jk} \\ &= aa_k^* = \left(\sum_j a_j e_j \right) a_k^* \\ &= \left(\sum_j a_j a_k^* \right) e_j. \end{aligned}$$

□

The projections E_a are special cases of the so-called **ket-bra-operators** (see e.g. [GVF01, p 71]). These (and their symbolic Dirac notation) are defined as

$$\begin{aligned} |r\rangle\langle s| : E &\longrightarrow F, \\ b &\longmapsto r(s|b), \end{aligned}$$

where E and F are Hilbert modules over a C^* -algebra $\mathcal{A}$, $r \in F$ and $s \in E$. For $E = F$, we have $E_a = |a\rangle\langle a|$. There are some relations among the ket-bra-operators:

$$\begin{aligned} r(s|ba) &= r(s|b)a, \\ |r\rangle\langle s|^* &= |s\rangle\langle r|, \\ |r\rangle\langle s| \circ |t\rangle\langle u| &= |r(t|s)\rangle\langle u| = |r\rangle\langle u(s|t)| \end{aligned}$$

hence the finite sums of ket-bra-operators from E to E form a self-adjoint algebra $End_{\mathcal{A}}^{00}(E)$ contained in $End_{\mathcal{A}}(E)$. The operators in $End_{\mathcal{A}}^{00}(E)$ are called operators **of $\mathcal{A}$ -finite rank**. The norm closure $End_{\mathcal{A}}^0(E)$ of $End_{\mathcal{A}}^{00}(E)$ contains the so-called **$\mathcal{A}$ -compact operators**. Clearly, $End_{\mathcal{A}}^0(\mathcal{A}^n) = \mathbb{M}_n(\mathcal{A})$ holds.

4.2. The modules $a\mathcal{A}$. We now turn to the examination of the modules $a\mathcal{A}$ onto which the E_a project. This subsection is quite technical. After showing how to “normalize” an arbitrary $a \in \mathcal{A}^n \setminus \{0\}$ to $\tilde{a}$ such that $(\tilde{a}|\tilde{a})$ and $E_{\tilde{a}}$ are projections with $a\mathcal{A} = \tilde{a}\mathcal{A}$, we define the *support* $S(M)$ of a projective submodule $M \subseteq \mathcal{A}^n$ and show that there is an $a \in M$ such that $S(a) = S(M)$ if M is finitely generated. This is used in the proof that the finitely generated projective submodules of $\mathcal{A}^n$ for which $End_{\mathcal{A}}(M)$ is *abelian* are exactly those of the form $a\mathcal{A} = \tilde{a}\mathcal{A}$. To show this, we also need the fact that $End_{\mathcal{A}}^0(P\mathcal{A}^n) = PM_n(\mathcal{A})P$ holds (Lemma 2.18 in [GVF01]).

All this is a preparation for the following subsection, where quasipoints of $\mathcal{R} \simeq \mathbb{M}_n(\mathcal{A})$ are regarded as families of projective submodules of $\mathcal{A}^n$ rather than families of projections.

Lemma 19. *Let $a \in \mathcal{A}^n$. $a\mathcal{A}$ is a closed submodule of $\mathcal{A}^n$.*

Proof. Let $(a\alpha_n)_{n \in \mathbb{N}}$ be a sequence in $a\mathcal{A}$ converging to $b \in \mathcal{A}^n$. As before, let $supp a := \bigcup_{k \leq n} supp a_k$, then $supp a = supp(a|a)$. Without loss of generality, one can assume that $\alpha_n(\beta) = 0$ holds for $\beta \neq supp a$, since the sequence $(a\alpha_n)$ remains unchanged by that. From

$$|a\alpha_n - a\alpha_m|^2 = |\alpha_n - \alpha_m|^2(a|a)$$

for all $n, m \in \mathbb{N}$ it follows that $(\alpha_n)_{n \in \mathbb{N}}$ is a Cauchy sequence in $\mathcal{A}$. Thus $(\alpha_n)_{n \in \mathbb{N}}$ converges to some $\alpha \in \mathcal{A}$ and we get

$$|a\alpha_n - a\alpha|^2 = |\alpha_n - \alpha|^2(a|a) \longrightarrow 0 \quad \text{for } n \longrightarrow \infty,$$

that is, $b = a\alpha \in a\mathcal{A}$. □

Let $M \subseteq \mathcal{A}^n$ be some submodule. The **orthogonal complement** $M^\perp$ of M is given by (see [WeO93, p 248])

$$M^\perp := \{b \in \mathcal{A}^n \mid \forall a \in M : (b|a) = 0\}.$$

For $M = a\mathcal{A}$, we obtain

$$\begin{aligned} (a\mathcal{A})^\perp &= \{b \in \mathcal{A}^n \mid \forall \alpha \in \mathcal{A} : (b|a\alpha) = 0\} \\ &= \{b \in \mathcal{A}^n \mid \forall \alpha \in \mathcal{A} : (b|a)\alpha = 0\} \\ &= \{b \in \mathcal{A}^n \mid (b|a) = 0\}. \end{aligned}$$

Obviously, $M \cap M^\perp = 0$ holds for any submodule M . A submodule M is called **complementable** if $M \oplus M^\perp = \mathcal{A}^n$ holds. One can show that M is complementable if and only if it is the image of some projection (Cor. 15.3.9 in [WeO93]).

Subsequently, it will be demonstrated how to normalize an arbitrary $a \in \mathcal{A}^n \setminus \{0\}$ to $\tilde{a}$ such that $(\tilde{a}|\tilde{a})$ is a projection in $\mathcal{A}$ and $E_{\tilde{a}}$ is the projection in $\mathbb{M}_n(\mathcal{A})$ onto $a\mathcal{A} = \tilde{a}\mathcal{A}$. Let $a \in \mathcal{A}^n \setminus \{0\}$ be such that $(a|a)$ is not a projection. For $n \in \mathbb{N}$, let

$$A_n := \{\beta \in \Omega \mid (a|a)(\beta) > \frac{1}{n}\}.$$

A_n is open and $\overline{A_n}$ is open and closed, since Ω is extremely disconnected. For an appropriate $n_0 \in \mathbb{N}$, $A_n \neq \emptyset$ holds for all $n \geq n_0$.

Let $\alpha_n \in \mathcal{A}$ be defined by

$$\alpha_n(\beta) := \begin{cases} (a|a)(\beta)^{-\frac{1}{2}} & \text{for } \beta \in \overline{A_n} \\ 0 & \text{for } \beta \notin \overline{A_n} \end{cases}.$$

Then $a\alpha_n \in a\mathcal{A}$ and

$$(a\alpha_n|a\alpha_n) = (a|a)\alpha_n^2 = \begin{cases} 1 & \text{on } \overline{A_n} \\ 0 & \text{on } \Omega \setminus \overline{A_n} \end{cases},$$

therefore $(a\alpha_n|a\alpha_n)$ is a projection in $\mathcal{A}$. Let $E_n := E_{a\alpha_n}$ be the projection onto $a\alpha_n\mathcal{A}$ given by $a\alpha_n$. According to the definition of α_n , $\alpha_n\mathcal{A}$ is the closed ideal

$$\alpha_n\mathcal{A} = \{\alpha \in \mathcal{A} \mid \text{supp } \alpha \subseteq \overline{A_n}\} = \chi_{\overline{A_n}}\mathcal{A}.$$

For these ideals it holds that

$$\forall n \in \mathbb{N} : \alpha_n\mathcal{A} \subseteq \alpha_{n+1}\mathcal{A}.$$

Moreover, with $S(a) := \text{supp } a \subseteq \Omega = \mathcal{Q}(\mathcal{A})$, it holds that $\overline{\bigcup_n \alpha_n\mathcal{A}} = \mathcal{A}_{\chi_{S(a)}}$: the inclusion “ $\subseteq$ ” is clear from $\overline{A_n} \subseteq S(a)$. Let $b \in \mathcal{A}^n$ be such that $S(b) \subseteq S(a)$, with no loss of generality $b \geq 0$. Then $(b\chi_{\overline{A_n}})_{n \geq n_0}$ is increasing monotonously. One gets

$$\sup_{n \in \mathbb{N}} b\chi_{\overline{A_n}} = b \sup_{n \in \mathbb{N}} \chi_{\overline{A_n}}.$$

From

$$\bigcup_n A_n = \{\beta \in \Omega \mid (a|a)(\beta) > 0\},$$

we have

$$\overline{\bigcup_n A_n} = \overline{\bigcup_n A_n} = S(a)$$

and thus $\sup_n \chi_{\overline{A_n}} = \chi_{S(a)}$, therefore

$$b = \sup_n b \chi_{\overline{A_n}}.$$

According to Dini's theorem, $b \chi_{\overline{A_n}}$ converges uniformly to b , and we have shown that $\overline{\bigcup_n \alpha_n \mathcal{A}} = \mathcal{A} \chi_{S(a)}$ holds.

The central carrier of the projection E_n is $\chi_{\overline{A_n}}$. $E_n \leq E_{n+1}$ holds, since for all $b \in \mathcal{A}^n$,

$$\begin{aligned} E_n E_{n+1} b &= a \alpha_n (a \alpha_n | E_{a \alpha_{n+1}} b) = a \alpha_n (a \alpha_n | a \alpha_{n+1}) (a \alpha_{n+1} | b) \\ &= a \alpha_n (a \alpha_{n+1} | a \alpha_{n+1}) (a \alpha_n | b) = a \alpha_n \chi_{\overline{A_{n+1}}} (a \alpha_n | b) \\ &= a \alpha_n (a \alpha_n | b) = E_n b, \end{aligned}$$

where we used $S(a \alpha_n) = \overline{A_n} \subseteq \overline{A_{n+1}}$ in the penultimate step. Let $E := \bigvee_{n \in \mathbb{N}} E_n$. The image of E_n is $a \mathcal{A} \chi_{\overline{A_n}}$, so

$$\text{im } E = a \mathcal{A} \chi_{S(a)} = a \mathcal{A}.$$

When defining E , one cannot simply assume the properties of the Hilbert space situation.

The sesquilinear form

$$\begin{aligned} \mathcal{A}^n \times \mathcal{A}^n &\longrightarrow \mathcal{A}, \\ (b, c) &\longmapsto (E_n b | c) \end{aligned}$$

can be written as

$$\begin{aligned} (E_n b | c) &= (a \alpha_n (a \alpha_n | b) | c) = \alpha_n^* (a | c) (a \alpha_n | b)^* \\ &= \alpha_n^* (a | c) (b | a) \alpha_n = \chi_{\overline{A_n}} (a | a)^{-\frac{1}{2}} | c) (b | a | a)^{-\frac{1}{2}}, \end{aligned}$$

and for $n \rightarrow \infty$, the right hand side converges to

$$(E b | c) := \chi_{S(a)} (a | a)^{-\frac{1}{2}} | b) (a | a)^{-\frac{1}{2}} | c).$$

Here we use the fact that every bounded continuous function f on an open dense subset $G \subseteq \Omega$ of the Stone space Ω can be extended to a continuous function $\tilde{f}$ on the whole of Ω (see Cor. III.1.8 in [TakI02]): the support $S(a) = \text{supp } a \subseteq \Omega$ is open and closed, $G(a) := \{\beta \in \Omega \mid a(\beta) \neq 0\}$ is open and dense in $S(a)$ and the mapping $a(a|a)^{-\frac{1}{2}} : G(a) \rightarrow \mathbb{C}^n$ is continuous und bounded, therefore it can be extended to a continuous mapping $S(a) \rightarrow \mathbb{C}^n$, which will also be denoted by $a(a|a)^{-\frac{1}{2}}$. Define

$$\tilde{a}(\beta) := \begin{cases} a(a|a)^{-\frac{1}{2}} & \text{for } \beta \in S(a) \\ 0 & \text{for } \beta \in \Omega \setminus S(a) \end{cases}$$

and from this, $E_{\tilde{a}}$. Then it holds for all $b, c \in \mathcal{A}^n$ that

$$(E_{\tilde{a}}b|c) = (\tilde{a}|c)(\tilde{a}|b) = \chi_{S(a)}(a(a|a)^{-\frac{1}{2}}|c)(a(a|a)^{-\frac{1}{2}}|b).$$

Obviously, $E_{\tilde{a}}$ is the same as the limit E of the projections E_n defined above. We showed both ways of the definition, because we will need the sets A_n for the following lemma.

Let $a \in \mathcal{A}^n \setminus \{0\}$, and let $\tilde{a} \in \mathcal{A}^n$ be as defined above. Then

$$(\tilde{a}|\tilde{a}) = (a|a)^{-1}(a|a) = 1$$

holds on $G(a)$ and hence also on $S(a)$. Thus $(\tilde{a}|\tilde{a})$ is a projection in $\mathcal{A}$.

Lemma 20. *For the closed submodules, we have $\tilde{a}\mathcal{A} = a\mathcal{A}$.*

Proof. From $\overline{\bigcup_n \mathcal{A}\chi_{A_n}} = \mathcal{A}\chi_{S(a)}$, one gets $a\mathcal{A} = \overline{\bigcup_n a\mathcal{A}\chi_{A_n}}$. Moreover, it holds that

$$a\mathcal{A}\chi_{A_n} = a(a|a)^{-\frac{1}{2}}\mathcal{A}\chi_{A_n} = \tilde{a}\mathcal{A}\chi_{A_n}.$$

It follows from this and $S(a) = S(\tilde{a})$ that

$$a\mathcal{A} = \overline{\bigcup_n a\mathcal{A}\chi_{A_n}} = \overline{\bigcup_n \tilde{a}\mathcal{A}\chi_{A_n}} = \tilde{a}\mathcal{A}.$$

□

Of course, $\tilde{a} = a$ if $(a|a)$ is a projection in $\mathcal{A}$, so one obtains

Proposition 21. *For every $a \in \mathcal{A}^n$, $E_{\tilde{a}}$ is an abelian projection with image $a\mathcal{A}$.*

$E_{\tilde{a}}$ is the unique projection from $\mathcal{A}^n$ onto $a\mathcal{A}$: let $Q : \mathcal{A}^n \rightarrow a\mathcal{A}$ be a projection. Then

$$\forall c \in (a\mathcal{A})^\perp : (Qc|Qc) = (Qc|c) = 0,$$

so $Q|_{(a\mathcal{A})^\perp} = 0$. Let $Qa = a\alpha$. From $Q^2 = Q$, $a\alpha^2 = a\alpha$, therefore $\alpha(\beta) \in \{0, 1\}$ on $S(a)$. Since $Q\mathcal{A}^n = a\mathcal{A}$, it follows that $\alpha = 1$ holds on $S(a)$, so $Qa = a$. Let $b \in \mathcal{A}^n$, $b = a\gamma + a'$ be such that $\gamma \in \mathcal{A}$ and $a' \in (a\mathcal{A})^\perp$. Such a decomposition exists, since $a\mathcal{A}$ is the image of a projection and hence a complementable submodule of $\mathcal{A}^n$. We have $Qb = a\gamma$ and

$$\begin{aligned} E_{\tilde{a}}b &= \tilde{a}(\tilde{a}|b) = \tilde{a}(\tilde{a}|a)\gamma = \tilde{a}(\tilde{a}|(a|a)^{\frac{1}{2}}\tilde{a})\gamma \\ &= \tilde{a}(a|a)^{\frac{1}{2}}\gamma = a\gamma = Qb, \end{aligned}$$

so $Q = E_{\tilde{a}}$.

Next, we will define the support of a submodule $M \subseteq \mathcal{A}^n$. For this, we will need a

Remark 22. *Let $U, V \subseteq \Omega$ be open. If $U \cap V = \emptyset$, then $\overline{U} \cap \overline{V} = \emptyset$.*

Proof. Let $\overline{U} \cap \overline{V} \neq \emptyset$ and $\beta \in \overline{U} \cap \overline{V}$. Since $\overline{U}$ is open, $\overline{U} \cap V \neq \emptyset$. Since V is open, $U \cap V \neq \emptyset$. □

Let M be a *projective submodule* of $\mathcal{A}^n$, i.e. $M = P\mathcal{A}^n$ for some projection $P \in \mathbb{M}_n(\mathcal{A})$ (see p 89 in [GVF01]). Then M is finitely generated.

$$\text{ann}(M) := \{\alpha \in \mathcal{A} \mid M\alpha = 0\},$$

the **annihilator of M** , is a closed ideal in $\mathcal{A}$. Let M be finitely generated, $\{g_1, \dots, g_r\} \subseteq M$ a system of generators. For $\alpha \in \mathcal{A}$, let $P(\alpha) := \{\beta \in \Omega \mid \alpha(\beta) \neq 0\}$ and for $a \in \mathcal{A}^n$ let $P(a) := \{\omega \in \Omega \mid a(\omega) \neq 0\}$. Then

$$\alpha a = 0 \iff P(\alpha) \cap P(a) = \emptyset,$$

and thus, according to the above remark,

$$\alpha a = 0 \iff S(\alpha) \cap S(a) = \emptyset.$$

Let

$$P(M) := \{\beta \in \Omega \mid \exists a \in M : \beta \in P(a)\}.$$

$S(M) := \overline{P(M)}$ is called the **support of M** . Since $\{g_1, \dots, g_r\}$ is a system of generators of M , $P(M) = \bigcup_{k \leq r} P(g_k)$ holds and hence $S(M) = \bigcup_{k \leq r} S(g_k) = \bigcup_{a \in M} S(a)$. The set $S(M)$ is open and closed. $\alpha \in \text{ann}(M)$ holds if and only if $\alpha g_k = 0$ for all $k \leq r$, therefore

$$\begin{aligned} \alpha &\in \text{ann}(M) \\ \iff S(\alpha) \cap S(g_k) &= \emptyset \quad (k \leq r) \\ \iff S(\alpha) \cap S(M) &= \emptyset \\ \iff S(\alpha) &\subseteq \Omega \setminus S(M) \\ \iff \alpha|_{S(M)} &= 0. \end{aligned}$$

This shows:

Remark 23. $\text{ann}(M)$ is the vanishing ideal of the closed-open set $S(M)$.

Lemma 24. $I_n \chi_{S(M)}$ is the central carrier of the projection $P_M \in \mathbb{M}_n(\mathcal{A})$ from $\mathcal{A}^n$ onto M .

Proof. Since $\chi_{S(M)} = 1$ on $S(a)$ for all $a \in M$, we have $\chi_{S(M)} a = a$ for all $a \in M$ and hence $\chi_{S(M)} P_M = P_M$, i.e. $C_{P(M)} \leq I_n \chi_{S(M)}$. Let $p \in \mathcal{A}$ be a projection such that $I_n p \geq P_M$. Then $pa = pP_M a = P_M a = a$ for all $a \in M$, so $p = 1$ on $S(a)$, that is, $p \geq \chi_{S(a)}$. It follows that $p \geq \chi_{S(M)}$. $\square$

Lemma 25. Let M be a finitely generated, projective submodule of $\mathcal{A}^n$, and let $a, b \in M$. Then there is a $c \in M$ such that $S(a) \cup S(b) \subseteq S(c)$.

Proof. Regard the decomposition $b = a\alpha + a'$ with $\alpha \in \mathcal{A}$ and $a' \in (a\mathcal{A})^\perp$. Such a decomposition always exists, since $a\mathcal{A}$ is a projective and hence complementable submodule of $\mathcal{A}^n$. Therefore each $b \in \mathcal{A}^n$ can be decomposed, in particular each $b \in M \subseteq \mathcal{A}^n$. We have $a' = b - a\alpha \in M$, so $a + a' \in M$ and

$$\begin{aligned} (b|b) &= \alpha^* \alpha (a|a) + (a'|a'), \\ (a + a'|a + a') &= (a|a) + (a'|a'), \end{aligned}$$

therefore

$$\begin{aligned} (a|a)(\beta) > 0 &\implies (a + a'|a + a')(\beta) > 0, \\ (b|b)(\beta) > 0 &\implies (a|a)(\beta) > 0 \text{ or } (a'|a')(\beta) > 0 \\ &\implies (a + a'|a + a')(\beta) > 0. \end{aligned}$$

It follows that $S(a|a) := \text{supp}(a|a) \subseteq S(a + a'|a + a')$ and $S(b|b) \subseteq S(a + a'|a + a')$, that is, $S(a) \cup S(b) \subseteq S(a + a')$. $\square$

Corollary 26. *Let M be as before. Then there is an $a \in M$ such that $S(a) = S(M)$. a can be chosen such that $(a|a)$ is a projection.*

Proof. Let $u_1, \dots, u_r \in M$ with $M = u_1\mathcal{A} + \dots + u_r\mathcal{A}$. According to the lemma above, there is an $a \in M$ such that $S(M) = \bigcup_{k \leq r} S(u_k) \subseteq S(a) \subseteq S(M)$, so $S(M) = S(a)$. Since $\tilde{a}\mathcal{A} = a\mathcal{A} \subseteq M$, $\tilde{a} \in M$, $(\tilde{a}|\tilde{a})$ is a projection and $S(\tilde{a}) = S(a)$. $\square$

Lemma 27. *For ket-bra-operators $|b\rangle\langle a|, |v\rangle\langle u| \in \mathbb{M}_n(\mathcal{A})$ it holds that $|b\rangle\langle a| \circ |v\rangle\langle u| = (a|v)|b\rangle\langle u|$ ($a, b, u, v \in M \subseteq \mathcal{A}^n$).*

Proof. Let $c \in M$. Then

$$\begin{aligned} |b\rangle\langle a| \circ |v\rangle\langle u|(c) &= b(a||v\rangle\langle u|(c)) = b(a|v(u|c)) \\ &= b(a|v)(u|c) = (a|v)|b\rangle\langle u|(c). \end{aligned}$$

$\square$

Let $M = P\mathcal{A}^n$ be a projective submodule. Lemma 2.18 in [GVF01] tells us that $\text{End}_{\mathcal{A}}^0(M) = P\mathbb{M}_n(\mathcal{A})P$, hence if P is abelian, so is $\text{End}_{\mathcal{A}}^0(P\mathcal{A}^n)$ and, in particular, $\text{End}_{\mathcal{A}}^{00}(P\mathcal{A}^n)$ is abelian. We will now characterize the finitely generated projective submodules $M \subseteq \mathcal{A}^n$ for which $\text{End}_{\mathcal{A}}^{00}(M)$, the set of $\mathcal{A}$ -linear mappings of $\mathcal{A}$ -finite rank from M to itself, is abelian. Let M be such a module. Let $u_1, \dots, u_r \in M$ be generators of M such that $(u_k|u_k) \in \mathcal{A}$ is a projection for all $k \leq r$. Moreover, let $a \in M$ be such that $(a|a)$ is a projection and $S(a) = S(M)$ holds. Then

$$|u_k\rangle\langle a| \circ |a\rangle\langle u_k| = (a|a)|u_k\rangle\langle u_k|$$

is a projection from M onto $(a|a)u_k\mathcal{A}$, and

$$|a\rangle\langle u_k| \circ |u_k\rangle\langle a| = (u_k|u_k)|a\rangle\langle a|$$

is a projection from M onto $(u_k|u_k)a\mathcal{A}$. Since $\text{End}_{\mathcal{A}}^{00}(M)$ is abelian by assumption, we get

$$\forall k \leq r : (a|a)u_k\mathcal{A} = (u_k|u_k)a\mathcal{A},$$

and since $S(a|a) = S(M)$, it holds that

$$\forall k \leq r : u_k\mathcal{A} = (u_k|u_k)a\mathcal{A} \subseteq a\mathcal{A}.$$

Thus it follows that

$$M = \sum_k u_k\mathcal{A} \subseteq a\mathcal{A} \subseteq M,$$

that is, $M = a\mathcal{A}$, so M is simply generated. Summing up, we have shown:

Proposition 28. *A projection $P \in \mathbb{M}_n(\mathcal{A}) = \text{End}_{\mathcal{A}}(\mathcal{A}^n)$ is abelian if and only if there is an $a \in \mathcal{A}^n$ such that $P = E_a$. Then $(a|a) \in \mathcal{A}$ is a projection and $I_n(a|a)$ is the central carrier of E_a . For $a \in \mathcal{A}^n$, $E_{\bar{a}}$ is the unique projection onto the simply generated submodule $a\mathcal{A}$ of $\mathcal{A}^n$. This submodule is projective. $a\mathcal{A}$ is free if and only if $S(a) = \Omega = \mathcal{Q}(\mathcal{A})$, i.e. if $E_{\bar{a}}$ has central carrier I_n .*

4.3. The equivalence relation on $\mathcal{A}^n$ and abelian quasipoints of $\mathcal{P}(\mathbb{M}_n(\mathcal{A}))$. In the following, we will regard quasipoints $\mathfrak{B}$ of $\mathcal{P}(\mathbb{M}_n(\mathcal{A}))$ as families of projective submodules of $\mathcal{A}^n$ rather than families of projections. The filter basis property of quasipoints will allow us to define a certain notion of germs on $\mathcal{A}$ and $\mathcal{A}^n$ and reduce the situation to that of finite dimensional vector spaces. The results known from that simple part of the theory (see Prop. 4) easily show that each quasipoint $\mathfrak{B} \in \mathcal{Q}(\mathbb{M}_n(\mathcal{A}))$ contains a submodule of the form $a_0\mathcal{A}$ and hence is abelian.

So let $\mathfrak{B} \in \mathcal{Q}(\mathbb{M}_n(\mathcal{A}))$ a quasipoint, regarded as a family of projective submodules M of $\mathcal{A}^n$. We have

$$P_{M \cap N} = P_M \wedge P_N,$$

if M, N as well as $M \cap N$ are algebraically finitely generated (see Sec. 15.4 of [WeO93]).

Let $P_M \wedge P_N$ be the minimum of the projections $P_M, P_N \in \mathbb{M}_n(\mathcal{A})$. Then

$$P_M \wedge P_N = P_K,$$

where K is the largest finitely generated closed submodule of $\mathcal{A}^n$ such that $K \subseteq M \cap N$ holds. We will use the notation $K = M \wedge N$.

As mentioned above, $I_n\mathcal{A}$ is the center of $\mathbb{M}_n(\mathcal{A})$.

$$\beta := \mathfrak{B} \cap I_n\mathcal{A}$$

is a quasipoint of $\mathcal{A}$, and it holds that

$$\beta \simeq \{\chi_{S(M)} \mid M \in \mathfrak{B}\}.$$

This can also be expressed in the following way:

$$\begin{aligned} \forall M \in \mathfrak{B} : \beta &\in \bigcup_{a \in M} S(a) \\ \iff \forall M \in \mathfrak{B} : \chi_{S(M)}(\beta) &= 1. \end{aligned}$$

Notice the double role of β : on the one hand as an element of the Stone spectrum (that is, the Gelfand spectrum) $\mathcal{Q}(\mathcal{A})$ of the center $I_n\mathcal{A} \simeq \mathcal{A}$ of $\mathbb{M}_n(\mathcal{A})$, on the other hand as a collection of characteristic functions $\chi_{S(M)}$ on this Stone spectrum.

We now define an equivalence relation on $\mathcal{A}^n$ that amounts to taking germs:

Definition 29. Let $n \in \mathbb{N}$. Two elements a, b of $\mathcal{A}^n$ are called **equivalent at the quasipoint** $\beta \in \mathcal{Q}(\mathcal{A})$, if there is a $p \in \beta$ such that $pa = pb$. Notation: $a \sim_\beta b$.

$\sim_\beta$ really is an equivalence relation: symmetry and reflexivity are obvious. Let $a \sim_\beta b, b \sim_\beta c$, then there are $p, q \in \beta$ such that $pa = pb, qb = qc \Rightarrow pq \in \beta$ and $pqa = pqb = pqc$, therefore $a \sim_\beta c$. Let $[a]_\beta$ be the equivalence class of $a \in \mathcal{A}^n$, and let $[\mathcal{A}^n]_\beta := \{[a]_\beta \mid a \in \mathcal{A}^n\}$.

Theorem 30. (i) $[\mathcal{A}]_\beta$ is a field,
(ii) $[\mathcal{A}^n]_\beta$ is an n -dimensional vector space over $[\mathcal{A}]_\beta$.

Proof. Let $a, b \in \mathcal{A}^n$. Then

$$[a]_\beta + [b]_\beta := [a + b]_\beta$$

is well defined, and also, for $\alpha \in \mathcal{A}$,

$$[a]_\beta [\alpha]_\beta := [a\alpha]_\beta$$

is well defined: from $a \sim_\beta a', b \sim_\beta b', \alpha \sim_\beta \alpha'$ it follows that there exist $p, q, r \in \beta$ such that

$$\begin{aligned} pa &= pa', qb = qb', r\alpha = r\alpha' \\ \implies pq(a+b) &= pqa + pqb = pqa' + pqb' = pq(a' + b') \end{aligned}$$

and

$$pr(a\alpha) = pa(r\alpha) = pa'(r\alpha') = pr(a'\alpha').$$

Furthermore,

$$[\alpha]_\beta [\gamma]_\beta := [\alpha\gamma]_\beta$$

defines a multiplication ($\alpha, \gamma \in \mathcal{A}$): it holds that

$$\begin{aligned} a \sim_\beta 0 &\iff \exists p \in \beta : pa = 0 \\ &\iff \exists p \in \beta : P(p) \cap P(a) = \emptyset \\ &\iff \exists p \in \beta : S(p) \cap S(a) = \emptyset \\ &\iff \beta \notin S(a). \end{aligned}$$

Let $\alpha \in \mathcal{A}$ be such that $[\alpha]_\beta \neq 0$. Then $\alpha^* \alpha \geq \varepsilon > 0$ holds on a closed-open neighbourhood W of β in $\Omega = \mathcal{Q}(\mathcal{A})$, since from $p\alpha \neq 0$ for all $p \in \beta$ and $p(\beta) = 1$, it follows that $\alpha(\omega) \neq 0$ holds on a neighbourhood of β . Thus $\chi_W \alpha$ is invertable on W and there is a $\gamma \in C(\Omega)$ such that $S(\gamma) = W$ and $\chi_W \alpha \gamma = \chi_W$, that is, $\alpha \gamma \sim_\beta 1$, so $[\alpha]_\beta [\gamma]_\beta = 1$. Since obviously the algebraic rules for multiplication and addition are fulfilled, it follows that $[\mathcal{A}]_\beta$ is a field and $[\mathcal{A}^n]_\beta$ is a vector space over $[\mathcal{A}]_\beta$. $([e_1]_\beta, \dots, [e_n]_\beta)$ is a basis of $[\mathcal{A}^n]_\beta$: let $a \in \mathcal{A}^n$, $a = \sum_{k=1}^n e_k a_k$, then $[a]_\beta = \sum_k [e_k]_\beta [a_k]_\beta$; if $\sum_k [e_k]_\beta [\gamma_k]_\beta = 0$, then there is some $p \in \beta$ such that

$$0 = p \sum_k e_k \gamma_k = \sum_k e_k (p \gamma_k),$$

so $p \gamma_k = 0$ for all k , that is, $[\gamma_k]_\beta = 0$ for all k . Thus we have

$$\dim_{[\mathcal{A}]_\beta} [\mathcal{A}^n]_\beta = n.$$

□

Let $M \subseteq \mathcal{A}^n$ be a submodule. Then $[M]_\beta := \{[\alpha]_\beta \mid \alpha \in M\}$ is a subspace of $[\mathcal{A}^n]_\beta$. If $N \subseteq \mathcal{A}^n$ is another submodule, then

Lemma 31. $[M \cap N]_\beta = [M]_\beta \cap [N]_\beta$.

Proof. The inclusion “ $\subseteq$ ” is trivial. Let $[a]_\beta \in [M]_\beta \cap [N]_\beta$. Then $[a]_\beta = [b]_\beta$ holds for some $b \in N$, so $pa = pb$ for a $p \in \beta$ and hence $pa \in M \cap N$. Since $p \sim_\beta 1$ in $\mathcal{A}$ (p is a projection), it follows that $[a]_\beta = [pa]_\beta \in [M \cap N]_\beta$. □

Corollary 32. $M, N \in \mathfrak{B} \implies [M \wedge N]_\beta \subseteq [M]_\beta \cap [N]_\beta$.

Proof. This follows from the lemma and $M \wedge N \subseteq M \cap N$. □

Let $M \in \mathfrak{B}$, then $[M]_\beta \neq 0$: assume that $[M]_\beta = 0$. Then

$$\begin{aligned} \forall a \in M \quad \exists p_a \in \beta : p_a a = 0 \\ \implies \forall a \in M : \beta \notin S(a), \end{aligned}$$

but $\beta \in S(M) = \bigcup_{a \in M} S(a)$, since $M \in \mathfrak{B}$. Thus we get

Remark 33. $\mathfrak{B}_\beta := \{[M]_\beta \mid M \in \mathfrak{B}\} \subseteq \mathbb{L}([\mathcal{A}^n]_\beta)$ is a filter basis in the lattice of subspaces of $[\mathcal{A}^n]_\beta$.

Proof. $0 \notin \mathfrak{B}_\beta$ and for $[M]_\beta, [N]_\beta \in \mathfrak{B}_\beta$, it holds that $[M]_\beta \cap [N]_\beta \supseteq [M \wedge N]_\beta \in \mathfrak{B}_\beta$. $\square$

Let $\tilde{\mathfrak{B}}_\beta$ be a quasipoint of $\mathbb{L}([\mathcal{A}^n]_\beta)$ containing $\mathfrak{B}_\beta$. There is exactly one line $[a_0]_\beta[\mathcal{A}]_\beta \in \tilde{\mathfrak{B}}_\beta$ such that

$$\forall M \in \mathfrak{B} : [a_0]_\beta[\mathcal{A}]_\beta \subseteq [M]_\beta,$$

because the discussion concerning atomic quasipoints for the lattice of subspaces of finite dimensional vector spaces holds (see Prop. 4). Of course we have $a_0 \approx_\beta 0$. It remains to show that $a_0\mathcal{A} \wedge M \neq 0$ holds for all $M \in \mathfrak{B}$. We have $P_{a_0\mathcal{A}} = E_{\tilde{a}_0}$ and, since $p_M \tilde{a}_0\mathcal{A} \subseteq M$ holds for some appropriate $p_M \in \beta$, we obtain

$$\begin{aligned} E_{p_M \tilde{a}_0} &\leq P_M \\ \iff p_M E_{\tilde{a}_0} &\leq P_M. \end{aligned}$$

From this, it follows (since $a_0 \approx 0$, we have $p_M \tilde{a}_0 \neq 0$ for all $p_M \in \beta$):

$$P_M \wedge E_{\tilde{a}_0} \geq p_M E_{\tilde{a}_0} \wedge E_{\tilde{a}_0} = p_M E_{\tilde{a}_0} > 0,$$

so $P_M \wedge E_{\tilde{a}_0} \neq 0$, that is, $a_0\mathcal{A} \wedge M \neq 0$. Since $\mathfrak{B}$ is a quasipoint, $a_0\mathcal{A} \in \mathfrak{B}$ holds from maximality. Summing up, we have shown:

Theorem 34. All quasipoints of $\mathbb{M}_n(\mathcal{A})$ are abelian.

The following remark clarifies the relation between several abelian projections in a single quasipoint:

Remark 35. Let $E_a, E_b \in \mathfrak{B}$ be abelian projections. There is some $r \in \mathcal{P}(\mathcal{A})$ such that $rE_a = rE_b$.

Proof. $E_c := E_a \wedge E_b$ is an abelian projection in $\mathfrak{B}$, so there are $p, q \in \mathcal{P}(\mathcal{A})$ such that $E_c = pE_a = qE_b$. Then $E_c = pE_a = p^2E_a = pqE_b$ holds and also $E_c = pqE_a$. If $p \notin \beta$, then $1 - p \in \beta$ and thus

$$0 = (1 - p)pE_a = (1 - p)E_c \in \mathfrak{B},$$

contradicting $0 \notin \mathfrak{B}$. Hence $p \in \beta$ and also $q \in \beta$, so $pq \in \beta$ and

$$pqE_a = pqE_b.$$

$\square$

If $\mathcal{R}$ is a type I_n algebra with trivial center, i.e. $\mathcal{R} \simeq \mathbb{M}_n(\mathbb{C}) \simeq \mathcal{L}(\mathbb{C}^n)$, then there are only atomic quasipoints (see Prop. 4 again). An atomic quasipoint is of the form

$$\mathfrak{B}_{\mathbb{C}x} := \{P \in \mathcal{P}(\mathcal{R}) \mid P_{\mathbb{C}x} \leq P\},$$

where $x \in \mathcal{H} \setminus \{0\}$. Of course, $P_{\mathbb{C}x}$ is an abelian projection, too. The Stone spectrum $\mathcal{Q}(\mathcal{R})$ is discrete in this case, since atomic quasipoints are isolated points of the Stone spectrum $\mathcal{Q}(\mathcal{R})$.

As a corollary of the results proved above, we obtain the main result:

Theorem 36. *Let $\mathcal{R}$ be a type I_n von Neumann algebra. The quasipoints of $\mathcal{A} = \mathcal{C}(\mathcal{R})$ parametrize the orbits of the unitary group $\mathcal{U}(\mathcal{R})$ acting on the Stone spectrum $\mathcal{Q}(\mathcal{R})$ of $\mathcal{R}$.*

Proof. All quasipoints of $\mathcal{R}$ are abelian (Thm. 34), i.e. we have $\mathcal{Q}(\mathcal{R}) = \mathcal{Q}^{ab}(\mathcal{A})$. Using this, we apply Thm. 11: the mapping

$$\begin{aligned} \zeta : \mathcal{Q}^{ab}(\mathcal{R}) &\longrightarrow \mathcal{Q}(\mathcal{A}), \\ \mathfrak{B} &\longmapsto \mathfrak{B} \cap \mathcal{A} \end{aligned}$$

is surjective. Since $\mathcal{R}$ is finite, we can replace partial isometries by unitary operators in Thm. 11. Therefore, $\zeta(\mathfrak{B}) = \zeta(\tilde{\mathfrak{B}})$ holds if and only if there is a unitary $U \in \mathcal{R}$ such that $U\mathfrak{B} = \tilde{\mathfrak{B}}$ (see Def. 12). It follows that the quasipoints of $\mathcal{A} = \mathcal{C}(\mathcal{R})$ parametrize the orbits of the unitary group $\mathcal{U}(\mathcal{R})$ acting on $\mathcal{Q}(\mathcal{R})$. $\square$

5. ACKNOWLEDGEMENT

I would like to thank H. de Groote for many helpful discussions and substantial support. Financial support by the Studienstiftung des Deutschen Volkes is gratefully acknowledged.

REFERENCES

- [Bir73] G. Birkhoff, *Lattice Theory (Revised Edition)* (Amer. Math. Soc., Providence 1973)
- [BirvNeu36] G. Birkhoff, J. von Neumann, “The logic of quantum mechanics”, *Ann. of Math.* **37** (1936), 823-843
- [Doe04] A. Döring, “Kochen-Specker theorem for von Neumann algebras”, <http://de.arXiv.org/abs/quant-ph/0408106>, accepted for publication in *Int. Journal of Theor. Phys.*
- [GVF01] J. M. Gracia-Bondia, J. C. Várilly, H. Figueroa, *Elements of Noncommutative Geometry* (Birkhäuser, Boston, Basel, Berlin 2001)
- [deG01] H. F. de Groote, “Quantum Sheaves”, preprint, <http://de.arxiv.org/abs/math-ph/0110035>
- [deG05] H. F. de Groote, *Observables*, Part I-IV (to appear 2005)
- [KadRinI97] R. V. Kadison, J. R. Ringrose, *Fundamentals of the Theory of Operator Algebras I* (AMS, Providence, Rhode Island 1997)
- [KadRinII97] R. V. Kadison, J. R. Ringrose, *Fundamentals of the Theory of Operator Algebras II* (AMS, Providence, Rhode Island 1997)
- [KocSpe67] S. Kochen, E. P. Specker, “The problem of hidden variables in quantum mechanics”, *Journal of Mathematics and Mechanics* **17** (1967), 59-87
- [Sto36] M. H. Stone, “The theory of representations for Boolean algebras”, *Trans. Amer. Math. Soc.* **40** (1936), 37-111
- [TakI02] M. Takesaki, *Theory of Operator Algebras I*, reprint of the 1979 edition (Springer, Berlin, Heidelberg, New York 2002)
- [WeO93] N. E. Wegge-Olsen, *K-Theory and C*-Algebras, A Friendly Approach* (Oxford University Press, Oxford, New York, Tokyo 1993)

E-mail address: adoering@math.uni-frankfurt.de